


Safety and Efficacy of the Combination of Selinexor, Daratumumab, and Dexamethasone (SDd) in Patients with Multiple Myeloma (MM) Previously Exposed to Proteasome Inhibitors and Immunomodulatory Drugs

Cristina Gasparetto, Suzanne Lentzsch, Gary Schiller, Natalie Callander, Sascha Tuchman, Christine Chen, Darrell White, Rami Kotb, Heather Sutherland, Michael Sebag, Muhamed Baljevic, William Bensinger, Richard LeBlanc, Chris Venner, Nizar Bahlis, Karla Rodriguez-Lopez, Heidi Sheehan, Jean-Richard Saint-Martin, Jatin Shah, Sharon Shacham, Michael Kauffman, Brea Lipe

Selinexor:

First-in-Class, Oral Selective Inhibitor of Nuclear Export (SINE)¹⁻⁴



Exportin 1 (XPO1) is the major nuclear export protein for:

- Tumor suppressor proteins (TSPs, e.g., p53, IκB, and FOXO)
- eIF4E-bound oncoprotein mRNAs (e.g., c-Myc, Bcl-xL, cyclins)
- Glucocorticoid receptor (GR)

XPO1 is overexpressed in MM:

- High **XPO1** levels enable cancer cells to escape TSP-mediated cell cycle arrest and apoptosis
- **XPO1** levels correlate with poor prognosis and drug resistance

Selinexor is an oral selective **XPO1 inhibitor**; preclinical data supports that selinexor:

- Reactivates multiple TSPs by preventing nuclear export
- Inhibits oncoprotein translation
- Reactivates GR signaling in presence of dexamethasone

Enhances daratumumab activity *in-vitro* against myeloma cells

Background / Rationale:

Daratumumab and Selinexor Activity in Heavily-Treated MM

SIRIUS: Daratumumab¹

Refractory to **PI** and **IMiD**

ORR: 29.2% (Overall)

ORR: 21.2% (Quad-Ref)

PFS: 3.7 months (Overall)

STORM: Selinexor + Dexamethasone²

Refractory to **Dara**, **PI**, and **IMiD**

ORR: 26.2%

ORR: 25.3% (Penta-Ref)

PFS: 3.7 months (Overall)

Both show single agent activity with ORR of 26-29% in RRMM

1. Lonial S, et al. Lancet. 2016;387:1551-60.

2. Chari A, et al. ASH 2018

STOMP Study Design

Primary Objective: Determine the maximum tolerated dose (MTD) and recommended Phase 2 dose (RP2D)

Patient Populations:

- **Arm SDd: selinexor + daratumumab + dexamethasone**
 - Patients who received ≥ 3 prior lines of therapy for MM, including a PI and an IMiD
 - Or patients with MM refractory to both a PI and an IMiD
- Arm SPd: selinexor + pomalidomide + dexamethasone (*EHA 2019 – Poster PF587*)
- Arm SVd: selinexor + bortezomib + dexamethasone
- Arm SKd: selinexor + carfilzomib + dexamethasone (*EHA 2019 – Poster PS1414*)
- Arm SRd: selinexor + lenalidomide + dexamethasone: >1 therapy and in patients with newly diagnosed MM

SDd Dosing Scheme: 3 + 3 design was used for dose escalation phase

Dose Level	Treatment Regimen	Patients Enrolled (Number of Patients with DLT)	Dose-Limiting Toxicity (DLT)
0	Selinexor, oral 60 mg (Days 1, 3) Twice Weekly Daratumumab, IV 16 mg/kg Once Weekly Dexamethasone, oral 20 mg Twice Weekly	3 (2)	Grade 2 fatigue and Grade 3 thrombocytopenia (both requiring reduction to 100 mg QW selinexor)
-1	Selinexor, oral 100 mg Once Weekly Daratumumab, IV 16 mg/kg Once Weekly Dexamethasone, oral 40 mg Once Weekly	6 (-)	No DLTs were reported in the 100 mg QW cohort

Patient Characteristics

Patient Characteristics	N
Enrolled as of May 1, 2019	34
60 mg selinexor BIW + 16 mg/kg daratumumab QW	3
100 mg selinexor QW + 16 mg/kg daratumumab QW (RP2D)	31
Median Age, Years (range)	68 (44 – 83)
Males : Females	19 (56%) : 15 (44%)
Median Time from Diagnosis to SDd Treatment, Years (range)	5.6 (<1 – 14)
Median Prior Regimens (range)	3 (2–10)
Proteasome Inhibitor (Treated: Refractory)	34 (100%) : 29 (85%)
Immunomodulatory Drug (Treated: Refractory)	34 (100%) : 26 (76%)
Autologous Stem Cell Transplant	24 (71%)
Daratumumab Treated	2 (6%)

Treatment-Related Non-Hematological Adverse Events in ≥10% Patients (RP2D Patients)

AE Term		100 mg Sel QW + 16 mg/kg Dara QW RP2D		
Gastrointestinal	Grade 1/2	Grade 3	Grade 4	Total (N=31)
Nausea	19 (61.3)	2 (6.5)	--	21 (67.7)
Dysgeusia	11 (35.5)	--	--	11 (35.5)
Diarrhea	9 (29.0)	1 (3.2)	--	10 (32.3)
Anorexia	10 (32.3)	--	--	10 (32.3)
Vomiting	8 (25.8)	--	--	8 (25.8)
Constipation	5 (16.1)	--	--	5 (16.1)
Constitutional				
Fatigue	13 (41.9)	5 (16.1)	--	18 (58.1)
Dizziness	5 (16.1)	--	--	5 (16.1)
Weight Loss	4 (12.9)	--	--	4 (12.9)
Other				
Hyponatremia	6 (19.4)	4 (12.9)	--	10 (32.3)
Insomnia	10 (32.3)	--	--	10 (32.3)
Vision Blurred	9 (29.0)	--	--	9 (29.0)
Hyperglycemia	5 (16.1)	1 (3.2)	--	6 (19.4)
Infusion Related Reaction	3 (9.7)	1 (3.2)	--	4 (12.9)

Safety data cutoff of May 1, 2019

Treatment-Related Hematological Adverse Events in $\geq 10\%$ Patients (RP2D Patients)

AE Term		100 mg Sel QW + 16 mg/kg Dara QW RP2D		
Hematologic	Grade 1/2	Grade 3	Grade 4	Total (N=31)
Thrombocytopenia	8 (25.8)	8 (25.8)	5 (16.1)	21 (67.7)
Anemia	9 (29.0)	9 (29.0)	--	18 (58.1)
Neutropenia	6 (19.4)	7 (22.6)	--	13 (41.9)
Leukopenia	4 (12.9)	8 (25.8)	--	12 (38.7)
Lymphopenia	1 (3.2)	3 (9.7)	1 (3.2)	5 (16.1)

- No treatment-related Grade 5 events were reported


Based on tolerability, the RP2D of SDd is selinexor 100 mg QW, daratumumab 16 mg/kg (per approved dosing) and dexamethasone 40 mg QW

Selinexor-Daratumumab-Dex: Efficacy


Category	N*	ORR (%)	VGPR (%)	PR [‡] (%)	CBR (%)	MR [‡] (%)	SD (%)	PD (%)
Daratumumab Naïve	30	22 (73%)	11 (37%)	11 (37%)	26 (87%)	4 (13%)	4 (13%)	--
All	32	22 (69%)	11 (34%)	11 (34%)	26 (81%)	4 (13%)	5 (16%)	1 (3%)

Responses were adjudicated according to the *International Myeloma Working Group* criteria,*two patients not evaluable for response withdrew consent prior to disease follow-up. ‡Two unconfirmed PRs, one unconfirmed MR. ORR=Overall Response Rate (VGPR+PR), VGPR=Very Good Partial Response, PR=Partial Response, MR=Minimal Response, SD=Stable Disease, PD=Progressive Disease, CBR=Clinical Benefit Rate (ORR+MR). Responses as of May 1, 2019 based on interim unaudited data.

SDd Efficacy by Prior Exposure : Dara-Naïve Patients (Efficacy Preserved Regardless of Prior Treatment)




SDd Efficacy – Patients with CBR ($\geq$ MR)




- Among patients with a response $\geq$ PR the median time on treatment was **7.7** months
- Median time to response was **1** month

SDd Efficacy – M-Protein Effect



Progression Free Survival



Conclusions – Safety & Efficacy

- **RP2D of SDd** – selinexor 100 mg, daratumumab 16 mg/kg and dexamethasone 40 mg, administered QW
- **Most Common G3/4 AEs** – thrombocytopenia, anemia, leukopenia, and neutropenia
- **Low-grade Gastrointestinal Side Effects** – common and expected, and can be managed with appropriate supportive care and/or dose modifications
- **ORR** – 73% in daratumumab-naïve patients
 - ~29% with daratumumab or 26% with selinexor alone
- **Clinical Benefit Rate** – 87% in daratumumab-naïve patients
- **Medians** – PFS and DOR medians have not been reached

Selinexor in combination with daratumumab/dexamethasone appears to be highly active, produces deep and durable responses in patients with RRMM, and warrants further investigation

Acknowledgments

Patients, their families, and caregivers

Investigators, co-investigators, and study teams at each participating center

- Duke University Cancer Center, Durham, North Carolina
- Columbia University, New York, NY
- UCLA Ronald Reagan Medical Center, Los Angeles, California
- Swedish Cancer Center, Seattle, WA
- Southern Alberta Cancer Research Institute, Calgary, Alberta
- Vancouver General Hospital, Vancouver, British Columbia
- Dalhousie University and QEII Health Sciences Center, Halifax, Nova Scotia
- Royal Victoria Hospital, Montreal, Québec
- Cancer Care Manitoba, Winnipeg, Manitoba
- Cross Cancer Institute, Edmonton, Alberta
- Hôpital Maisonneuve-Rosemont, Montreal, Québec
- Princess Margaret Cancer Center, Toronto, Ontario
- Myeloma Canada, Laval, Québec
- University of Rochester Medical College, New York, NY
- Moffitt Cancer Center, Tampa, FL
- University of Nebraska Medical Center

This study was sponsored by Karyopharm Therapeutics